


Reference List



Ying-Xiu Dai, Ying-Yen Chuang, Po-Yu Chen, and Chih-Chiang Chen.
Efficacy and safety of ablative resurfacing with a high-energy 1,064 nm Q-switched Nd:YAG laser for the treatment of facial acne scars in Asians.
Lasers Surg. Med. © 2019 Wiley Periodicals, Inc. (wileyonlinelibrary.com).
DOI 10.1002/lsm.23151

Guida S, Giovanni P, Bencini PL.
Picosecond laser treatment of atrophic and hypertrophic surgical scars: In vivo monitoring of results by means of 3D imaging and reflectance confocal microscopy.
Skin Res Technol. 2019;00:1–7. <https://doi.org/10.1111/srt.12755>

Haider Al-Sabak Muhsin A. Al-Dhalimi Wid Abbas Ali.
Treatment Of Verruca Vulgaris with Long Pulsed Nd:YAG Laser 1064 nm In Iraqi Patients.
KUFA JOURNAL FOR NURSING SCIENCES.VOL.9 No. 1 / 2019

Ploypailin Tantrapornpong, Punyaphat Sirithanabadeekul.
Efficacy and Safety of Using a Fractional Picosecond 1064-nm Laser for Treatment of Atrophic Acne Scars in Asian Skin Types.
TJPS Vol.42 (Supplement Issue) 2018

S. Guida, P.L. Bencini, G. Pellacani.
Picosecond laser for atrophic surgical scars treatment: in vivo monitoring of results by means of reflectance confocal microscopy.
Letter to the editor. *J EADV* 2018. DOI: 10.1111/jdv.15289

Nark-Kyoung Rho, Sung Tae Chung.
Treatment of a Full-thickness Laceration Scar on the Forehead Using a 1,064-nm Fractional Picosecond Laser and Polynucleotide Gel Injection.
Med Laser 2018;7(2):88-90. <https://doi.org/10.25289/ML.2018.7.2.88>

Variya Laothaworn & Premjit Juntongjin (2018):
Topical 3% tranexamic acid enhances the efficacy of 1064-nm Q-switched neodymium-doped yttrium aluminum garnet laser in the treatment of melasma.
Journal of Cosmetic and Laser Therapy, DOI:10.1080/14764172.2018.1427869

Murtadha Hashim Al-janabi, Noor Taha Ismaeel Ali, Karrar Deiaa Mohamed Al-Sabti, Muhsin A. Al-Dhalimi & Sahib Neamh Abdul Wahid (2017).
A new imaging technique for assessment of the effectiveness of long pulse Nd:YAG 532 nm laser in treatment of facial port wine stain.
Journal of Cosmetic and Laser Therapy, 19:7, 418-421, DOI:10.1080/14764172.2017.1341985

K. Greveling, E.P. Prens, N. ten Bosch and M. B. van Doorn.
Comparison of lidocaine/tetracaine cream and lidocaine/prilocaine cream for local anaesthesia during laser treatment of acne keloidalis nuchae and tattoo removal: results of two randomized controlled trials.
British Journal of Dermatology (2017) 176, pp81–86. DOI 10.1111/bjd.14848

Barakat MT, Moftah NH, El Khayyat MAM, and Abdelhakim ZA.
Significant reduction of inflammation and sebaceous glands size in acne vulgaris lesions after intense pulsed light treatment.
Dermatologic Therapy 2017;30: e12418. doi:10.1111/dth.12418.

El-Domyati M, Wael H, Noha M, Hamza A, Selwet M.
Hair follicle changes following intense pulsed light axillary hair reduction: histometrical, histological and immunohistochemical evaluation.
Archives of Dermatological Research (2017); 309(3):191– 200

Paradisi A, Ricci F, Sbrano P.
“Drug-resistant granuloma faciale”: treatment with carbon dioxide-GaAs laser.
Derm Ther 2016; 29:317–319.


Al-Dhalimi MA, Al-Janabi MH.
Split lesion randomized comparative study between long pulsed Nd:YAG laser 532 and 1,064 nm in treatment of facial port-wine stain.
Lasers Surg Med 2016;48:852–8.

Paolo Mezzana, Maurizio Valeriani & Roberto Valeriani (2016):
Combined fractional resurfacing (10600nm/1540nm): Tridimensional imaging evaluation of a new device for skin rejuvenation
Journal of Cosmetic and Laser Therapy, DOI:10.1080/14764172.2016.1202417

Paradisi A, Ricci F, Sbrano P.
“Drug-resistant granuloma faciale”: treatment with carbon dioxide-GaAs laser.
Dermatol Ther 2016; 29: 317–319.

Maluki AH, Al-Sabak HA (2015)
Split Face Comparative Study of using Fractional Erbium: Glass 1540 Nanometer and Nd: YAG Q-Switched 1064 Nanometer Laser Systems in the Treatment of Melasma.
J Dermatol Res Ther 1:002

Reference List



- Muhsin Al-Dhalimi & Akeel Jaber (2015)
Treatment of atrophic facial acne scars with fractional Er:Yag laser
Journal of Cosmetic and Laser Therapy, 17:4, 184-188, DOI: 10.3109/14764172.2015.1007067
- Mohammad Radmanesh & Zohreh Rafiei (2015)
Combination of CO2 and Q-switched Nd:YAG lasers is more effective than Q-switched Nd:YAG laser alone for eyebrow tattoo removal
Journal of Cosmetic and Laser Therapy, 17:2, 65-68, DOI:10.3109/14764172.2014.988724
- Leclère FM, Moreno-Moraga J, Alcolea JM, Casoli V, Mordon SR, Vogt PM, Trelles MA (2014)
Laser assisted lipolysis for neck and submental remodeling in Rohrich type I to III aging neck: a prospective study in 30 patients.
J Cosmet Laser Ther 16(6):284–289
- Muhsin A. Al-Dhalimi & Ali A. Abo Nasyria (2013)
A comparative study of the effectiveness of intense pulsed light wavelengths (650 nm vs 590 nm) in the treatment of striae distensae.
Journal of Cosmetic and Laser Therapy, 15:3, 120-125, DOI: 10.3109/14764172.2012.748200
- Al-Dhalimi MA, Abo Nasyria AA.
A comparative study of the effectiveness of intense pulsed light wavelengths (650 nm vs 590 nm) in the treatment of striae distensae.
J Cosmet Laser Ther 2013;15:120-5.
- Trelles MA, Mordon SR, Bonanad E, Moreno Moraga J, Heckmann A, Unglaub F, Betrouni N, Leclère FM (2013)
Laser-assisted lipolysis in the treatment of gynecomastia: a prospective study in 28 patients.
Lasers Med Sci 28:375–382
- Kassab AN, El Kharbotly A.
Management of ear lobule keloids using 980-nm diode laser.
Eur Arch Otorhinolaryngol. 2012;269(2):419–423.
- Leclère FM, Trelles M, Moreno-Moraga J, Servell P, Unglaub F, Mordon SR (2012)
980-nm laser lipolysis (LAL): About 674 procedures in 359 patients.
J Cosmet Laser Ther 14:67–73
- Moreira A, Leite I, Guedes R, et al.
Surgical treatment of rhinophyma using carbon dioxide (CO2) laser and pulsed dye laser (PDL).
J Cosmet Laser Ther 2010; 2010: 73–76.
- Madan V, Ferguson JE, August PJ.
Carbon dioxide laser treatment of rhinophyma: a review of 124 patients.
Br J Dermatol 2009; 161: 814–818.
- Bouzari N, Tabatabai H, Abbasi Z, et al.
Laser hair removal: comparison of long-pulsed Nd:YAG, long-pulsed alexandrite, and long-pulsed diode lasers.
Dermatol Surg 2004;30:498-502.
- Goon PK, Dalal M, Peart FC.
The gold standard for decortication of rhinophyma: combined erbium-YAG/CO2 laser.
Aesthetic Plast Surg 2004; 28: 456–460.
- Redett RJ, Manson PN, Goldberg N, et al.
Methods and results of rhinophyma treatment.
Plast Reconstr Surg 2001; 107: 1115–1123.